

Oncology Areas of Interest

(Investigator Initiated Studies and Research Collaborations)

R2810 (Cemiplimab)

Cemiplimab is available for dosing Q3W IV or Q6W IV

Types of proposals

- Combination with standard of care (SOC) including exploration of sequencing
- Studies in indications of high unmet need and unaddressed rare populations
- Novel study concepts investigating squamous histologies
- Studies incorporating ctDNA as an exploratory endpoint to generate early insights or as a key endpoint, treatment-guiding factor, or enrollment criterion in the neoadjuvant/ adjuvant/ perioperative setting (except in NMSC) may be considered on a very limited basis

NOT PREFERRED: Concomitant IO/RT

Areas of interest for NSCLC

- Stage I-III treatment settings
- Novel combinations

Areas of interest for non-melanoma skin cancer

- Novel combinations
- BCC
 - Combination studies in all treatment stages including post-anti-PD-1
 - Neoadjuvant setting (prior to surgery or RT)
 - High-risk populations
- CSCC
 - Treatment of high-risk pre-cancerous lesions and studies in post-IO settings including retreatment will be considered on a limited basis

Other tumor types

- SCLC, GI, genitourinary, and gynecological cancers will be considered on a limited basis

Note: Novel combinations must have rationale for complementary mechanisms of action using agents with established dose range and safety

(V6.0, May 2025)

Oncology Areas of Interest

(Investigator Initiated Studies and Research Collaborations)

REGN3767 (anti-LAG3)_fianlimab

Note: Fianlimab is dosed at 1600 mg IV Q3W in combination with cemiplimab

- The fianlimab program will consider only exceptional proposals on a very limited basis

(V7.0, Oct 2025)

REGN5093 (METxMET)_davutamig

Areas of interest for NSCLC

- Davutamig in patients with MET alterations who are unable to tolerate TKIs
- Combination studies of davutamig with TKIs (including EGFR-targeted or MET targeted therapies) in patients with MET-amplification or overexpression
- Combination studies of davutamig with cemiplimab (anti-PD-1) in MET alterations in immune-checkpoint inhibitor-naïve populations

Areas of interest in other tumor types

- Proposals addressing other indications where MET may be a driver or a mechanism of resistance to targeted therapy will be considered on a limited basis (i.e. CRC, GI cancers, mesothelioma, papillary RCC)

(v3.0, May 2025)